arXiv:1211.4572v1 [math.NT] 18 Nov 2012

SOME PROPERTIES ON THE INTEGRAL OF THE PRODUCT OF SEVERAL EULER POLYNOMIALS

TAEKYUN KIM

Abstract. In this paper, we study the formula for a product of two Euler polynomials. From this study, we derive some formulae for the integral of the product of two or more Euler polynomials.

1. Introduction

As is well known, the Bernoulli polynomials are given by the generating function as follows:

$$\frac{t}{e^t-1}e^{xt}=e^{B(x)t}=\sum_{n=0}^{\infty}B_n(x)\frac{t^n}{n!},\quad |t|<2\pi,$$

with the usual convention about replacing $B^n(x)$ by $B_n(x)$. In the special case $x=0$, $B_n(0)=B_n$ are called the n-th Bernoulli numbers (see [18-20]). The constants E_k in the Taylor series expansion

$$\frac{2}{e^t+1}=\sum_{n=0}^{\infty}E_n\frac{t^n}{n!},\quad |t|<\pi, \tag{1}$$

(cf. [2,11,12]) are known as the k-th Euler numbers. From the generating function of Euler numbers, we note that

$$E_0=1,\quad E_n=-\sum_{i=1}^{n}\binom{n}{i}E_i,\quad \text{for } n\in\mathbb{N}. \tag{2}$$

The first few are 1, $-\frac{1}{2}$, 0, $\frac{1}{4}$, ..., and $E_{2k}=0$ for $k=1,2,\ldots$. The Euler polynomials are defined by

$$\frac{2}{e^t+1}e^{xt}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!}=\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\binom{n}{k}E_kx^{n-k}\right)\frac{t^n}{n!}. \tag{3}$$

Thus, by (3), we get

$$E_n(x)=\sum_{k=0}^{n}\binom{n}{k}E_kx^{n-k}=\left(E+x\right)^n,\quad \text{(see [2,5,11,12])}, \tag{4}$$

with the usual convention about replacing E^n by E_n.

From (1), (2) and (4), we have

$$E_0=1,\quad E_n(1)+E_n=\left(E+1\right)^n+E_n=0,\quad \text{if}\quad n\geq 1. \tag{5}$$

By the definition of Euler polynomials, we easily see that

$$\sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!} = \frac{2}{e^t+1}e^{xt} = \frac{2}{1+e^{-t}}e^{-t(1-x)} = \sum_{n=0}^{\infty}(-1)^n E_n(1-x)\frac{t^n}{n!}. \tag{6}$$

From (6), we have the reflection symmetric relation for Euler polynomials as follows:

$$E_n(x) = (-1)^n E_n(1-x), \quad \text{(see [1-16])}. \tag{7}$$

By (4), we get

$$\begin{aligned} \frac{d}{dx}E_n(x) &= \sum_{l=0}^{n-1}\binom{n}{l}(n-l)x^{n-l-1}E_l = n\sum_{l=0}^{n-1}\binom{n-1}{l}E_l x^{n-1-l} \\ &= n\big(E+x\big)^{n-1} = nE_{n-1}(x). \end{aligned} \tag{8}$$

Thus, by (8), we see that

$$\begin{aligned} \int_0^1 E_n(x)dx &= \frac{1}{n+1}\int_0^1 \frac{d}{dx}E_{n+1}(x)dx \\ &= \frac{1}{n+1}\big(E_{n+1}(1)-E_{n+1}\big) = -\frac{2}{n+1}E_{n+1}, \quad \text{for} \quad n \in \mathbb{Z}_+ = \mathbb{N}\cup\{0\}. \end{aligned} \tag{9}$$

The gamma and beta functions are defined as the following definite integrals ($\alpha > 0$, $\beta > 0$):

$$\Gamma(\alpha) = \int_0^\infty e^{-t}t^{\alpha-1}dt, \quad \text{(see [20])}, \tag{10}$$

and

$$\mathrm{B}(\alpha,\beta) = \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}dt = \int_0^\infty \frac{t^{\alpha-1}}{(1+t)^{\alpha+\beta}}dt. \tag{11}$$

By (10) and (11), we get the following equation:

$$\Gamma(\alpha+1) = \alpha\Gamma(\alpha), \quad \mathrm{B}(\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad \text{(see [20])}. \tag{12}$$

In this paper we give some interesting properties of several Euler polynomials to express the integral of those polynomials from 0 to 1 in terms of beta and gamma functions. Finally, we derive some identities on the integral of the product of Euler polynomials.

2. On the integral of the product of Euler polynomials

Let us consider the integral for the product of Euler polynomials and x^n as follows:

$$\int_0^1 y^n E_n(x+y)dy = \sum_{l=0}^{n}\binom{n}{l}E_{n-l}(x)\int_0^1 y^{n+l}dy = \sum_{l=0}^{n}\frac{\binom{n}{l}E_{n-l}(x)}{n+l+1}. \tag{13}$$

On the other hand, by (7), we get

$$\begin{aligned}
\int_0^1 y^n E_n(x+y)dy &= (-1)^n \int_0^1 y^n E_n(1-(x+y))dy \\
&= (-1)^n \sum_{l=0}^{n} \binom{n}{l} E_{n-l}(-x) \int_0^1 y^n (1-y)^l dy \\
&= \sum_{l=0}^{n} \binom{n}{l} (-1)^l E_{n-l}(1+x) \mathrm{B}(n+1, l+1) \\
&= \sum_{l=0}^{n} \binom{n}{l} (-1)^l E_{n-l}(1+x) \frac{\Gamma(n+1)\Gamma(l+1)}{\Gamma(n+l+2)}.
\end{aligned} \tag{14}$$

Therefore, by (13) and (14), we obtain the following theorem.

Theorem 1. *For $n \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$, we have*

$$\sum_{l=0}^{n} \frac{\binom{n}{l} E_{n-l}(x)}{n+l+1} = \sum_{l=0}^{n} (-1)^l \left(\frac{E_{n-l}(1+x)}{n+l+1} \right) \frac{\binom{n}{l}}{\binom{n+l}{l}}.$$

In particular, $x = 0$,

$$\sum_{l=0}^{n} \frac{\binom{n}{l} E_{n-l}}{n+l+1} = (-1)^n \sum_{l=0}^{n} \frac{E_{n-l}}{n+l+1} \frac{\binom{n}{l}}{\binom{n+l}{l}}.$$

Let $n \in \mathbb{N}$ with $n \geq 3$. Then, by (9), we see that

$$\begin{aligned}
&\int_0^1 y^n E_n(x+y)dy = \frac{E_n(x+1)}{n+1} - \frac{n}{n+1} \int_0^1 y^{n+1} E_{n-1}(x+y)dy \\
&= \frac{E_n(x+1)}{n+1} - \frac{E_{n-1}(x+1)}{n+1} \frac{n}{n+2} + (-1)^2 \frac{n(n-1)}{(n+1)(n+2)} \int_0^1 y^{n+2} E_{n-2}(x+y)dy \\
&= \frac{E_n(x+1)}{n+1} - \frac{nE_{n-1}(x+1)}{(n+1)(n+2)} + (-1)^2 \frac{n(n-1)E_{n-2}(x+1)}{(n+1)(n+2)(n+3)} \\
&\qquad + (-1)^3 \frac{n(n-1)(n-2)}{(n+1)(n+2)(n+3)} \int_0^1 y^{n+3} E_{n-3}(x+y)dy.
\end{aligned} \tag{15}$$

Continuing this process, we obtain the following equation:

$$
\begin{aligned}
&\int_0^1 y^n E_n(x+y)dy \\
&= \frac{E_n(x+1)}{n+1} + \sum_{l=2}^{n-1} \frac{n(n-1)\cdots(n-l+2)(-1)^{l-1}}{(n+1)(n+2)\cdots(n+l)} E_{n-l+1}(1+x) \\
&\qquad + (-1)^{n-1}\frac{n(n-1)(n-2)\cdots 2}{n(n+1)\cdots(2n-1)}\int_0^1 y^{2n-1}E_1(x+y)dy \\
&= \frac{E_n(x+1)}{n+1} - \sum_{l=2}^{n-1} \frac{n(n-1)\cdots(n-l+2)(-1)^{l-1}}{(n+1)(n+2)\cdots(n+l)} E_{n-l+1}(1+x) \\
&\qquad + (-1)^{n-1}\frac{n!}{(n+1)(n+2)\cdots 2n}\Big(E_1(x+1) - \frac{1}{2n+1}\Big).
\end{aligned}
\tag{16}
$$

Therefore, by (14) and (16), we obtain the following theorem.

Theorem 2. *For $n \in \mathbb{N}$ with $n \geq 3$, we have*

$$
\begin{aligned}
\sum_{l=0}^{n}(-1)^l \frac{E_{n-l}(x+1)}{n+l+1}\frac{\binom{n}{l}}{\binom{n+l}{l}} &= \frac{E_n(x+1)}{n+1} + \sum_{l=2}^{n-1}\frac{\binom{n}{l}(n-l+2)(-1)^{l-1}}{\binom{n+l}{l}}E_{n-l+1}(1+x) \\
&\quad + \frac{(-1)^{n-1}}{\binom{2n}{n}}\Big(E_1(x+1) - \frac{1}{2n+1}\Big).
\end{aligned}
$$

In the special case, $x = 0$,

$$
\sum_{l=0}^{n}\frac{E_{n-l}}{n+l+1}\frac{\binom{n}{l}}{\binom{n+l}{l}} = \frac{E_n}{n+1} + \sum_{l=2}^{n-1}\frac{\binom{n}{l}(n-l+2)}{\binom{n+l}{l}}E_{n-l+1} + \frac{1}{\binom{2n}{n}}\Big(\frac{1}{2} + \frac{1}{2n+1}\Big).
$$

For $n \in \mathbb{N}$, we have

$$
\int_0^1 y^n E_n(x+y)dy = \frac{E_n(x)}{n+1} - \frac{n}{n+1}\int_0^1 y^{n-1}E_{n+1}(x+y)dy. \tag{17}
$$

By (17), we get

$$
\int_0^1 y^n E_n(x+y)dy = \frac{E_n(x)}{n+1} - \frac{n}{n+1}\sum_{l=0}^{n+1}\binom{n+1}{l}E_{n+1-l}(-x)(-1)^{n+1}\mathrm{B}(n, l+1). \tag{18}
$$

Therefore, by (14) and (18), we obtain the following theorem.

Theorem 3. *For $n \in \mathbb{N}$, we have*

$$
\frac{E_{n+1}(x)}{n+1} = \frac{1}{n+1}\sum_{l=0}^{n+1}\frac{\binom{n+1}{l}}{\binom{n+l}{l}}E_{n+1-l}(1+x)(-1)^l - \sum_{l=0}^{n}\frac{E_{n-l}(1+x)}{n+l+1}(-1)^l\frac{\binom{n}{l}}{\binom{n+l}{l}}
$$

In particular, $x = 0$,

$$
-\frac{E_{n+1}}{n+1} = \frac{1}{n+1}\sum_{l=0}^{n+1}\frac{\binom{n+1}{l}}{\binom{n+l}{l}}E_{n+1-l} + \sum_{l=0}^{n}\frac{\binom{n}{l}}{\binom{n+l}{l}}\frac{E_{n-l}}{n+l+1}.
$$

A definite integral for the multiplication of two Euler polynomials can be given by the following relation:

$$\begin{aligned}\int_0^1 E_n(x)E_m(x)dx &= \sum_{l=0}^{n}\binom{n}{l}E_l(-1)^m\sum_{k=0}^{m}\binom{m}{k}E_k\int_0^1 x^{n-l}(1-x)^{m-k}dx \\ &= \sum_{l=0}^{n}\sum_{k=0}^{m}\binom{n}{l}\binom{m}{k}(-1)^m E_lE_k\mathrm{B}(n-l+1,m-k+1).\end{aligned} \tag{19}$$

Let $m,n\in\mathbb{N}_+$ with $m\geq 1$. Then we see that

$$\begin{aligned}\int_0^1 E_n(x)E_m(x)dx &= -\frac{m}{n+1}\int_0^1 E_{n+1}(x)E_{m-1}(x)dx \\ &= (-1)^2\frac{m(m-1)}{(n+1)(n+2)}\int_0^1 E_{n+2}(x)E_{m-2}(x)dx.\end{aligned} \tag{20}$$

Continuing this process, we have

$$\int_0^1 E_n(x)E_m(x)dx = (-1)^{m-1}\frac{m(m-1)\cdots 2}{(n+1)(n+2)\cdots(n+m-1)}\int_0^1 E_{n+m-1}(x)E_1(x)dx. \tag{21}$$

It is easy to show that

$$\int_0^1 E_{n+m-1}(x)E_1(x)dx = -\frac{1}{m+n}\int_0^1 E_{m+n}(x) = \frac{2E_{n+m-1}}{(m+n)(m+n+1)}. \tag{22}$$

Thus, by (20), (21) and (22), we get

$$\begin{aligned}\int_0^1 E_n(x)E_m(x)dx &= (-1)^{m+1}\frac{2m!E_{n+m+1}}{(n+1)(n+2)\cdots(n+m)(n+m+1)} \\ &= 2(-1)^{m+1}\frac{m!n!}{(n+m)!}\frac{E_{n+m+1}}{n+m+1}.\end{aligned} \tag{23}$$

Therefore, by (19) and (23), we obtain the following theorem.

Theorem 4. *For $m,n\in\mathbb{Z}_+$ with $m\geq 1$, we have*

$$-2\frac{E_{n+m+1}}{\binom{n+m}{n}(n+m+1)} = \sum_{l=0}^{n}\sum_{k=0}^{m}\binom{n}{l}\binom{m}{k}E_lE_k\frac{\Gamma(n-l+1)\Gamma(m-k+1)}{\Gamma(n+m-l-k+2)}.$$

Moveover

$$E_{n+m+1} = -\frac{1}{2}\sum_{l=0}^{n}\sum_{k=0}^{m}\frac{\binom{n}{l}\binom{m}{k}\binom{n+m}{n}}{\binom{n+m-l-k}{n-l}}\left(\frac{(n+m+1)E_lE_k}{n+m-l-k+1}\right).$$

From (3), we note that

(24)

$$\begin{aligned}
&\sum_{m,n=0}^{\infty}\Big(mE_{m-1}(x)E_n(x)+nE_m(x)E_{n-1}(x)\Big)\frac{t^m}{m!}\frac{s^n}{n!}\\
&\qquad=\frac{d}{dx}\left(\frac{4e^{(s+t)x}}{(e^t+1)(e^s+1)}\right)=(s+t)\frac{4}{(e^t+1)(e^s+1)}e^{(s+t)x}\\
&\qquad=\left(\frac{s+t}{e^{(s+t)-1}}e^{(s+t)x}\right)\left(4-\frac{4}{e^t+1}-\frac{4}{e^s+1}\right)\\
&\qquad=\left(\sum_{l=0}^{\infty}B_l(x)\frac{(s+t)^l}{l!}\right)\left(4-2\sum_{r=0}^{\infty}E_r\frac{t^r}{r!}-2\sum_{r=0}^{\infty}E_r\frac{s^r}{r!}\right)\\
&\qquad=\left(\sum_{m,n=0}^{\infty}B_{m+n}(x)\frac{t^ms^n}{m!n!}\right)\left(-2\sum_{r=0}^{\infty}\frac{E_{2r+1}}{(2r+1)!}\Big(t^{2r+1}+s^{2r+1}\Big)\right)\\
&\qquad=\sum_{m,n=0}^{\infty}\left(-2\sum_{r=0}^{\infty}\frac{E_{2r+1}}{(2r+1)!}\frac{B_{m+n}(x)}{m!n!}\Big(t^{2r+m+1}s^n+s^{2r+1+n}t^m\Big)\right)\\
&\qquad=\sum_{m,n=0}^{\infty}\left(-2\sum_{r=0}^{\infty}\frac{E_{2r+1}}{(2r+1)!}\Big(B_{m-2r-1+n}(x)\frac{t^ms^nm!}{(m-2r-1)!n!m!}\Big)\right.\\
&\qquad\qquad\qquad\left.+\frac{B_{m+n-2r-1}(x)}{(n-2r-1)!m!}s^nt^m\frac{n!}{n!}\right)\\
&\qquad=\sum_{m,n=0}^{\infty}\left(-2\sum_{r=0}^{\infty}E_{2r+1}B_{m-2r-1+n}(x)\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)\right)\frac{t^ms^n}{m!n!}.
\end{aligned}$$

By comparing coefficients on the both sides in (24), we obtain the following theorem.

Theorem 5. *For $m,n\in\mathbb{N}$, we have*

$$mE_{m-1}(x)E_n(x)+nE_m(x)E_{n-1}(x)=-2\sum_{r=0}^{\infty}E_{2r+1}B_{m-2r-1+n}(x)\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right).$$

From (8) and (9), we note that

$$\begin{aligned}
\frac{d}{dx}\Big(E_m(x)E_n(x)\Big)&=mE_{m-1}(x)E_n(x)+nE_m(x)E_{n-1}(x)\\
&=-2\sum_{r=0}^{\infty}E_{2r+1}B_{m-2r-1+n}(x)\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right).
\end{aligned}\tag{25}$$

Thus, by (25), we get

$$\begin{aligned}
E_m(x)E_n(x)&=\int\frac{d}{dx}\Big(E_m(x)E_n(x)\Big)dx\\
&=-2\sum_{r=0}^{\infty}E_{2r+1}\frac{B_{m+n-2r}(x)}{m+n-2r}\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)+C,
\end{aligned}\tag{26}$$

where C is some constant.

For $m+n\geq 2$, we have

$$\int_0^1 E_m(x)E_n(x)dx = C. \tag{27}$$

By (23) and (27), we get

$$C = 2(-1)^{m+1}\frac{m!n!}{(n+m)!}\frac{E_{n+m+1}}{n+m+1} = (-1)^{m+1}\frac{2}{\binom{n+m}{n}}\frac{E_{n+m+1}}{n+m+1}. \tag{28}$$

Therefore, by (26) and (28), we obtain the following theorem.

Theorem 6. *For $m, n \in \mathbb{N}$ with $m+n\geq 2$, we have*

$$\begin{aligned}&E_m(x)E_n(x)\\ &= -2\sum_{r=0}^{\infty}\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)\frac{E_{2r+1}B_{m+n-2r}(x)}{m+n-2r} + 2(-1)^{m+1}\frac{E_{n+m+1}}{\binom{n+m}{n}(n+m+1)}.\end{aligned}$$

Note that

$$\begin{aligned}&\int_0^1 B_{m+n-2r}(x)E_p(x)dx = \left[\frac{B_{m+n-2r+1}(x)E_p(x)}{m+n-2r+1}\right]_0^1 - p\int_0^1 \frac{B_{m+n-2r+1}(x)E_{p-1}(x)}{m+n-2r+1}dx\\ &= -2\frac{B_{m+n-2r+1}E_p}{m+n-2r+1} - p\left[\frac{B_{m+n-2r+2}(x)E_{p-1}(x)}{(m+n-2r+1)(m+n-2r+2)}\right]_0^1\\ &\qquad + p(p-1)(-1)^2\int_0^1 \frac{B_{m+n-2r+2}(x)E_{p-2}(x)}{(m+n-2r+1)(m+n-2r+2)}dx.\end{aligned}$$

Continuing this process, we obtain

$$\begin{aligned}&\int_0^1 B_{m+n-2r}(x)E_p(x)dx\\ &= 2p!\sum_{l=0}^{p-1}\frac{B_{m+n-2r+l}E_{p-l+1}(-1)^l}{(m+n-2r+1)\cdots(m+n-2r+l)} + \frac{p!(-1)^{p-1}\int_0^1 B_{m+n-2r+p-1}(x)E_1(x)dx}{(m+n-2r+1)\cdots(m+n-2r+p-1)}\\ &= 2p!\sum_{l=0}^{p-1}\frac{B_{m+n-2r+l}E_{p-l+1}(-1)^l}{(m+n-2r+1)\cdots(m+n-2r+l)} + \frac{2p!(-1)^pB_{m+n-2r+p}E_1}{(m+n-2r+1)\cdots(m+n-2r+p)}\\ &= 2p!\sum_{l=0}^{p}\frac{B_{m+n-2r+l}E_{p-l+1}(-1)^l}{(m+n-2r+1)\cdots(m+n-2r+l)} = 2p!\sum_{l=0}^{p}\frac{B_{m+n-2r+l}E_{p-l+1}}{\binom{n+m-2r+l}{l}}\frac{(-1)^l}{l!}.\end{aligned} \tag{29}$$

From Theorem 6, we can derive the following equation (30). For $m, n, p \in \mathbb{N}$, we have

$$\begin{aligned}
&\int_0^1 E_m(x)E_n(x)E_p(x)dx = \int_0^1 E_p(x)(E_m(x)E_n(x))dx \\
&= -2\sum_{r=0}^{\infty}\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)\frac{E_{2r+1}}{m+n-2r}\int_0^1 B_{m+n-2r}(x)E_p(x)dx \\
&\quad + 2(-1)^{m+1}\frac{E_{n+m+1}}{\binom{n+m}{n}(n+m+1)}\int_0^1 E_p(x)dx.
\end{aligned} \tag{30}$$

By (29) and (30), we get

$$\begin{aligned}
&\int_0^1 E_m(x)E_n(x)E_p(x)dx \\
&= -4p!\sum_{r=0}^{\infty}\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)\frac{E_{2r+1}}{n+m-2r}\sum_{l=1}^{p}\frac{B_{m+n-2r+l}E_{p-l+1}}{\binom{n+m-2r+l}{l}}\frac{(-1)^l}{l!} \\
&\quad + 4\frac{(-1)^m E_{n+m+1}E_{p+1}}{\binom{n+m}{n}(n+m+1)(p+1)}.
\end{aligned} \tag{31}$$

On the other hand,

$$\begin{aligned}
&\int_0^1 E_m(x)E_n(x)E_p(x)dx \\
&= (-1)^p\sum_{l=0}^{m}\sum_{j=0}^{n}\sum_{k=0}^{p}\binom{m}{l}\binom{n}{j}\binom{p}{k}E_{m-l}E_{n-j}E_k\int_0^1 x^{l+j}(1-x)^{p-k}dx \\
&= (-1)^p\sum_{l=0}^{m}\sum_{j=0}^{n}\sum_{k=0}^{p}\binom{m}{l}\binom{n}{j}\binom{p}{k}E_{m-l}E_{n-j}E_k \mathrm{B}(l+j+1, p-k+1) \\
&= (-1)^p\sum_{l=0}^{m}\sum_{j=0}^{n}\sum_{k=0}^{p}\binom{m}{l}\binom{n}{j}\binom{p}{k}E_{m-l}E_{n-j}E_k\frac{(l+j)!(p-k)!}{(l+j+p-k+1)!}.
\end{aligned} \tag{32}$$

Therefore, by (31) and (32), we obtain the following equation:

$$\begin{aligned}
&-4p!\sum_{r=0}^{\infty}\left(\binom{m}{2r+1}+\binom{n}{2r+1}\right)\frac{E_{2r+1}}{n+m-2r}\sum_{l=1}^{p}\frac{B_{m+n-2r+l}E_{p-l+1}}{\binom{n+m-2r+l}{l}}\frac{(-1)^l}{l!} \\
&+4\frac{(-1)^m E_{n+m+1}E_{p+1}}{\binom{n+m}{n}(n+m+1)(p+1)} = (-1)^p\sum_{l=0}^{m}\sum_{j=0}^{n}\sum_{k=0}^{p}\frac{\binom{m}{l}\binom{n}{j}\binom{p}{k}E_{m-l}E_{n-j}E_k}{\binom{l+j+p-k}{l+j}(l+j+p-k+1)},
\end{aligned}$$

where $m, n, p \in \mathbb{N}$.

ACKNOWLEDGEMENTS. The author would like to express his sincere gratitude to referees for their valuable comments and suggestions.

References

[1] M. Abramowitz, I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, 55 (1964).

[2] S. Araci, D. Erdal, J. Seo, *A study on the fermionic p-adic q-integral on $\mathbb{Z}_p$ associated with weighted q-Bernstein q-Genocchi polynomials*, Abstract and Applied Analysis, 2011(2011), Article ID **649248**, 10 pages.

[3] A. Bayad, J. Chikhi, *Non linear recurrences for Apostol-Bernoulli-Euler number of higher order*, Adv. Stud. Contemp. Math. **22**(2012), no.1, 1–6.

[4] A. Bayad, T. Kim, B. Lee, S.-H. Rim, *Some identities on the Bernstein polynomials associated with q-Euler polynomials*, Abstract and Applied Analysis, 2011(2011), Article ID **294715**, 10 pages.

[5] L. Carlitz, *Note on the integral of the product of several Bernoulli polynomials*, J. Londen Math. Soc. 34 (1959), 361–363.

[6] M. Cenkci, Y. Simsek, V. Kurt, *Multiple two-variable q-L-function and its behavior at $s = 0$*, Russ. J. Math. Phys. 15 (2008), 447–459.

[7] J. Choi, D. S. Kim, T. Kim, Y.-H. Kim, *A note on some identities of Frobenius-Euler numbers and polynomials*, Int. J. Math. Math. Sci. 2012(2012), Article ID 861797, 7 pages.

[8] T. Kim, *Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. 16 (2009), 484–491.

[9] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. 16 (2009), 93–96.

[10] T. Kim, *A note on q-Bernstein polynomials*, Russ. J. Math. Phys. 18 (2011), 73–82.

[11] T. Kim, *A note on the Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **17**(2008), 131–136.

[12] T. Kim, *Some identities on the q-integral representation of the product of several q-Bernstein-type polynomials*, Abstract and Applied Analysis, **2011**(2011), Article ID 634675, 11 pages.

[13] H. Ozden, I. N. Cangul, Y. Simsek, *Remarks on q-Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), 41–48.

[14] H. Ozden, I. N. Cangul, Y. Simsek, *Multivariate interpolation functions of higher-order q-Euler numbers and their applications*, Abstr. Appl. Anal. 2008(2008), Art. ID. **390857**, 16 pages.

[15] C. S. Ryoo, *Some identities of the twisted q-Euler numbers and polynomials associated with q-Bernstein polynomials*, Proc. Jangjeon Math. Soc. **14** (2011), 239–348.

[16] C. S. Ryoo, *Some relations between twisted q-Euler numbers and Bernstein polynomials*, Adv. Stud. Contemp. Math. **21**(2011), 217–223.

[17] Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. **16** (2008), 251–278.

[18] Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Adv. Stud. Contemp. Math. **11** (2005), 205–218.

[19] Y. Simsek, *Special functions related to Dedekind-type DC-sums and their applications*, Russ. J. Math. Phys. 17 (2010), 495–508.

[20] D. G. Zill, M. R. Cullen, *Advanced Engineering Mathematics*, Jones and Bartlett Publishers, Inc., 2006.

Taekyun Kim

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail: tkkim@kw.ac.kr